

Branching ratio for $B \rightarrow K_1\gamma$ decay in next-to-leading order in LEET

M. Jamil Aslam

*National Centre for Physics and Department of Physics,
Quaid-i-Azam University,
Islamabad, Pakistan*

Riazuddin

*National Centre for Physics,
Quaid-i-Azam University,
Islamabad, Pakistan*

Branching ratio for $B \rightarrow K_1\gamma$ at next-to-leading order of α_s has been calculated in Large Energy Effective Theory. By incorporating the higher twist effects in light cone decay amplitude for axial K meson, it is shown that the form factor is not sensitive to these twists.

I. INTRODUCTION

Rare B decays involving flavor-changing-neutral-current (FCNC) transitions, such as $b \rightarrow s\gamma$, have received a lot of theoretical interest [1]. First measurements of the decay $B \rightarrow X_s\gamma$ were reported by the CLEO collaboration [2]. These decays are now being investigated more precisely in experiments at the B factories. The current world average based on the improved measurements by the CLEO [3], ALEPH [4] and BELLE collaborations, $\mathcal{B}(B \rightarrow X_s\gamma) = (3.22 \pm 0.40) \times 10^{-4}$, is in good agreement with the estimates of the standard model (SM) [5–7], which we shall take as $\mathcal{B}(B \rightarrow X_s\gamma) = (3.50 \pm 0.50) \times 10^{-4}$, reflecting the parametric uncertainties dominated by the scheme-dependence of the quark masses. The decay $B \rightarrow X_s\gamma$ also provides useful constraints on the parameters of the supersymmetric theories, which in the context of the minimal supersymmetric standard model (MSSM) have been recently updated [8].

Exclusive decays involving the $b \rightarrow s\gamma$ transition are best exemplified by the decay $B \rightarrow K^*\gamma$, which provide abundant issues for both theorists and experimentalists. After the first measurement at CLEO, $B \rightarrow K^*\gamma$ is now also measured in Belle and Babar:

$$\mathcal{B}(B^0 \rightarrow K^{*0}\gamma) = \begin{cases} (4.09 \pm 0.21 \pm 0.19) \times 10^{-5} & \text{Belle [9]} \\ (4.23 \pm 0.40 \pm 0.22) \times 10^{-5} & \text{BaBar [10]} \\ (4.55 \pm 0.70 \pm 0.34) \times 10^{-5} & \text{CLEO [11]} \end{cases} \quad (1)$$

$$\mathcal{B}(B^+ \rightarrow K^{*+}\gamma) = \begin{cases} (4.40 \pm 0.33 \pm 0.24) \times 10^{-5} & \text{Belle [9]} \\ (3.83 \pm 0.62 \pm 0.22) \times 10^{-5} & \text{BaBar [10]} \\ (3.76 \pm 0.86 \pm 0.28) \times 10^{-5} & \text{CLEO [11]} \end{cases} \quad (2)$$

On theoretical side there have been noticeable advances in $B \rightarrow K^*\gamma$ for a decade. QCD corrections at next-to-leading order (NLO) of $O(\alpha_s)$ have already been considered [12–14]. Relevant Wilson coefficients have been improved up to three loop level calculations [15,16]. Recent developments of the QCD factorizations helped one to calculate the hard spectator contributions systematically in the factorized form through the convolution at the heavy quark limit [17–19]. The detailed analysis of $B \rightarrow K^*\gamma$ has also been done at next to leading order in effective theories, such as large energy effective theory (LEET) [20], and in soft-collinear effective theory (SCET) [23].

In addition to K^* , higher resonances of kaon also deserve much attention. Recently, Belle has announced the first measurement of $B \rightarrow K_1^+(1270)\gamma$ [24]

$$\mathcal{B}(B^+ \rightarrow K_1^+\gamma) = (4.28 \pm 0.94 \pm 0.43) \times 10^{-5} \quad (3)$$

Among many reasons to focus on the higher resonances, the first one is that they share lot of the things with $B \rightarrow K^*\gamma$, like at quark level both of them are governed by $b \rightarrow s\gamma$. Therefore all the achievements of $b \rightarrow s\gamma$ can be used in these decays, e.g. the same operators in the operator product expansion and the same Wilson coefficients that are available. The light cone distribution amplitudes (DA) are same except the overall factor of γ_5 and this gives few differences in many calculations [25]. Secondly, it was suggested that $B \rightarrow K_{\text{res}}(\rightarrow K\pi\pi)\gamma$ can provide a direct measurement of the photon polarization [26] and it was shown that large polarization asymmetry $\approx 33\%$ has been produced due to decay of B meson through the kaon resonances. In the presences of anomalous right-handed couplings, the polarization can be severely reduced in the parameter space allowed by current experimental bounds of $B \rightarrow X_s\gamma$. It was also

argued that the B factories can now make a lot of $B\bar{B}$ pairs, enough to check the anomalous couplings through the measurement of the photon polarization.

The theorists are also facing challenges from the discrepancy between their predictions and experiments. It was pointed out that the form factor obtained using the LEET approach for $B \rightarrow K^*\gamma$ is found to be smaller compared to the values obtained by QCD sum rules or light-cone sum rules (LCSR) [20]. At this stage, the source of this mismatch is not well understood.

On $B \rightarrow K_1\gamma$ side the situation is more complicated. Based on the QCDF framework combined with the LCSR results, it is predicted that $\mathcal{B}(B^0 \rightarrow K_1^0(1270)\gamma) = (0.828 \pm 0.335) \times 10^{-5}$ at the NLO of α_s which is very small as compared to the experimental value [cf. Eq. (3)] [25]. The value of the relevant form factor has been extracted from the experimental data and its value is found to be $F_+^{K_1(1270)}(0) = 0.32 \pm 0.03$ which is very large as compared to $F_+^{K_1(1270)}(0)|_{\text{LCSR}} = 0.14 \pm 0.03$ obtained by the LCSR. These are contrary to the case of $B \rightarrow K^*\gamma$ where the form factor obtained from LCSR is larger than the LEET one and the source of discrepancy is not yet known. But for $B \rightarrow K_1\gamma$ case the possible candidates to explain this discrepancy have also been discussed in detail in the literature [28].

In this paper the branching ratio for $B \rightarrow K_1\gamma$ at NLO of α_s are calculated using the LEET approach [29,30]. We follow the same frame work as done by Ali et. al. [20] for $B \rightarrow K^*\gamma$, because $B \rightarrow K_1\gamma$ shares many things with it. The only difference is the DA for the daughter meson. As K_1 is an axial vector and is distinguished by vector by the γ_5 in the gamma structure of DA and some non perturbative parameters. But the presence of γ_5 does not alter the calculation, give the same result for the perturbative part. The higher twist terms are also included through the Gegenbauer moments in the Gegenbauer expansion. The calculation with out Gegenbauer has already been done in QCD factorization frame work and using the LCSR results for form factors and decay constant [25,28].

At next-to-leading order of α_s there are the contributions from the operators O_2 O_7 and O_8 which will be discussed in detail. The paper is organized as follows. In Section II we give the short flavor of the leading order calculation for $B \rightarrow K_1\gamma$ decay process. The section III deals with the hard spectator contributions in $B \rightarrow K_1\gamma$ decays while in section IV $O(\alpha_s)$ corrected matrix element for above mentioned decays is discussed in detail. The resulting branching ratio and related discussion appear in sec. V. The concluding remarks are given at the end.

II. LEADING ORDER CONTRIBUTIONS

The effective Hamiltonian for $b \rightarrow s\gamma$ can be written as

$$\mathcal{H}_{\text{eff}}(b \rightarrow s\gamma) = -\frac{G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum_{i=1}^8 C_i(\mu) O_i(\mu) , \quad (4)$$

where

$$\begin{aligned} O_1 &= (\bar{s}_i c_j)_{V-A} (\bar{c}_j b_i)_{V-A} , \\ O_2 &= (\bar{s}_i c_i)_{V-A} (\bar{c}_j b_j)_{V-A} , \\ O_3 &= (\bar{s}_i b_i)_{V-A} \sum_q (\bar{q}_j q_j)_{V-A} , \\ O_4 &= (\bar{s}_i b_j)_{V-A} \sum_q (\bar{q}_j q_i)_{V-A} , \\ O_5 &= (\bar{s}_i b_i)_{V-A} \sum_q (\bar{q}_j q_j)_{V+A} , \\ O_6 &= (\bar{s}_i b_j)_{V-A} \sum_q (\bar{q}_j q_i)_{V+A} , \\ O_7 &= \frac{em_b}{8\pi^2} \bar{s}_i \sigma^{\mu\nu} (1 + \gamma_5) b_i F_{\mu\nu} , \\ O_8 &= \frac{g_s m_b}{8\pi^2} \bar{s}_i \sigma^{\mu\nu} (1 + \gamma_5) T_{ij}^a b_j G_{\mu\nu}^a . \end{aligned} \quad (5)$$

Here i, j are color indices, and we neglect the CKM element $V_{ub} V_{us}^*$ as well as the s -quark mass. The leading contribution to $B \rightarrow K_1\gamma$ comes from the electromagnetic operator O_7 as shown in Fig. a. As in the case of the real photon emission ($q^2 = 0$), the only form factor appears in the calculation is $\xi_{\perp}^{(K_1)}$. Therefore one can write

$$\begin{aligned}\langle O_7 \rangle_A &\equiv \langle K_1(p', \epsilon) \gamma(q, e) | O_7 | B(p) \rangle \\ &= \frac{em_b}{4\pi^2} \xi_{\perp}^{(K_1)} \left[\epsilon^* \cdot q(p + p') \cdot e^* - \epsilon^* \cdot e^* (p^2 - p'^2) + i\epsilon_{\mu\nu\alpha\beta} \epsilon^{*\mu} \epsilon^{*\nu} q^\alpha (p + p')^\beta \right],\end{aligned}\quad (6)$$

with $\epsilon^{*\nu}$ and e^μ being the polarization vector for axial kaon and the photon respectively. The decay rate is straightforwardly obtained to be [25]

$$\Gamma(B \rightarrow K_1 \gamma) = \frac{G_F^2 \alpha m_b^2 m_B^3}{32\pi^4} |V_{tb} V_{ts}^*|^2 \left(1 - \frac{m^2}{m_B^2}\right)^3 |\xi_{\perp}^{(K_1)}|^2 |C_7^{\text{eff}(0)}|^2, \quad (7)$$

where α is the fine-structure constant and $C_7^{\text{eff}(0)}$ is the effective Wilson coefficient at leading order.

III. MATRIX ELEMENTS AT NEXT-TO-LEADING ORDER OF $O(\alpha_s)$

At next to leading order of α_s , there are the contributions from the operators O_2 and O_8 along with that of the O_7 in $B \rightarrow K_1 \gamma$ decay. Each operator has its vertex contribution and hard spectator contribution terms which we calculate explicitly.

A. Hard Spectator Contribution

The Hard spectator contribution is well described by the convolution between the hard kernel T_k and the light cone distribution amplitudes of the involved mesons, Φ_B and Φ_{K_1} and can be written as $\Phi_B \otimes T_k \otimes \Phi_{K_1}$. The corresponding decay amplitude can be calculated in the form of convolution formula, whose leading term can be expressed as [20]

$$\Delta \mathcal{M}^{(\text{HSA})} = \frac{4\pi\alpha_s C_F}{N_c} \int_0^1 du \int_0^\infty dl_+ M_{jk}^{(B)} M_{li}^{(\rho)} \mathcal{T}_{ijkl} \quad (8)$$

where N_c is the number of colors, $C_F = (N_c^2 - 1)/(2N_c)$ is the Casimir operator eigenvalue in the fundamental representation of the color $SU(N_c)$ group. The leading-twist two-particle light-cone projection operators $M_{jk}^{(B)}$ [31,33] and $M_{li}^{(K_1)}$ [32,33] of B - and K_1 -mesons in the momentum representation are:

$$M_{jk}^{(B)} = -\frac{if_B M}{4} \left[\frac{1+\psi}{2} \left\{ \phi_+^{(B)}(l_+) \not{l}_+ + \phi_-^{(B)}(l_+) \left(\not{l}_- - l_+ \gamma_\perp^\mu \frac{\partial}{\partial l_\perp^\mu} \right) \right\} \gamma_5 \right]_{jk} \Big|_{l=(l_+/2)n_+}, \quad (9)$$

$$M_{li}^{(K_1)} = -\frac{i}{4} \left[f_\perp^{(K_1)} (\not{\epsilon}^* \not{p}) \gamma_5 \phi_\perp^{(K_1)}(u) + f_\parallel^{(K_1)} \left(\not{p} \frac{m}{E} (v \varepsilon^*) \right) \gamma_5 \phi_\parallel^{(K_1)}(u) \right]_{li}, \quad (10)$$

where f_B is the B -meson decay constant, $f_\parallel^{(K_1)}$ and $f_\perp^{(K_1)}$ are the longitudinal and transverse K_1 -meson decay constants, respectively, and ε_μ is the K_1 -meson polarization vector. These projectors include also the leading-twist distribution amplitudes $\phi_+^{(B)}(l_+)$ and $\phi_-^{(B)}(l_+)$ of the B -meson and $\phi_\parallel^{(K_1)}(u)$ and $\phi_\perp^{(K_1)}(u)$ of the K_1 -meson. $\mathcal{T}_{ijkl}$ is the hard-scattering amplitude. The Kinematical relations are used to calculate the hard spectator contributions are [33]

$$\begin{aligned}p_b^\mu &\simeq m_b v^\mu, & l^\mu &= \frac{l_+}{2} n_+^\mu + l_\perp^\mu + \frac{l_-}{2} n_-^\mu \\ k_1^\mu &\simeq u E n_-^\mu + k_\perp^\mu + O(k_\perp^2), & k_2^\mu &\simeq \bar{u} E n_-^\mu - k_\perp^\mu + O(k_\perp^2), \\ v^2 &= 1, & v^\mu &= (n_-^\mu + n_+^\mu)/2 & E &\simeq M/2 \\ q^\mu &= \omega n_+^\mu & \omega &= M/2\end{aligned}$$

To calculate $\mathcal{T}_{ijkl}$ let's consider the contribution from all the possible diagrams as done for the $B \rightarrow V \gamma$ [20].

1. Spectator corrections due to the electromagnetic dipole operator O_7

The corresponding diagrams are presented in Fig. 1 and the explicit expression is given by

$$\begin{aligned} \mathcal{T}_{ijkl}^{(1)} = & -i \frac{G_F}{\sqrt{2}} V_{td}^* V_{tb} C_7^{\text{eff}}(\mu) \frac{em_b(\mu)}{4\pi^2} \frac{[\gamma_\mu]_{kl}}{(l-k_2)^2} \\ & \times \left[(q\sigma e^*)(1+\gamma_5) \frac{\not{p}_b + \not{l} - \not{k}_2 + m_b}{(p_b + l - k_2)^2 - m_b^2} \gamma_\mu \right. \\ & \left. + \gamma_\mu \frac{\not{k}_1 + \not{k}_2 - \not{l}}{(k_1 + k_2 + l)^2} (q\sigma e^*)(1+\gamma_5) \right]_{ij} \end{aligned} \quad (11)$$

where the short hand notation is used for $(q\sigma e^*) = \sigma^{\mu\nu} q_\mu e_\nu^*$.

2. Spectator corrections due to the chromomagnetic dipole operator O_8

The corresponding diagrams are presented in Fig. 2. The first two diagrams (Fig. 2a) show the corrections for the case when the photon is emitted from the flavor changing quark line and the result is

$$\begin{aligned} \mathcal{T}_{ijkl}^{(2a)} = & -i \frac{G_F}{\sqrt{2}} V_{td}^* V_{tb} C_8^{\text{eff}}(\mu) \frac{em_b(\mu)}{4\pi^2} [\gamma_\nu]_{kl} \frac{(l-k_2)_\mu}{(l-k_2)^2} \\ & \times \left[\not{\epsilon}^* \frac{\not{p}_b + \not{l} - \not{k}_2}{(p_b + l - k_2)^2} \sigma_{\mu\nu} (1+\gamma_5) \right. \\ & \left. + \sigma_{\mu\nu} (1+\gamma_5) \frac{\not{k}_1 + \not{k}_2 - \not{l} + m_b}{(k_1 + k_2 + l)^2 - m_b^2} \not{\epsilon}^* \right]_{ij} \end{aligned} \quad (12)$$

Fig. 2b contains the diagrams with the photon emission from the spectator quark which results into the following hard-scattering amplitude:

$$\begin{aligned} \mathcal{T}_{ijkl}^{(2b)} = & i \frac{G_F}{\sqrt{2}} V_{td}^* V_{tb} C_8^{\text{eff}}(\mu) \frac{eQ_{d[u]}m_b(\mu)}{4\pi^2} \\ & \times [\sigma_{\mu\nu} (1+\gamma_5)]_{ij} \frac{(p_b - k_1)_\mu}{(p_b - k_1)^2} \\ & \times \left[\gamma_\nu \frac{\not{p}_b + \not{l} - \not{k}_1}{(p_b + l - k_1)^2} \not{\epsilon}^* + \not{\epsilon}^* \frac{\not{k}_1 + \not{k}_2 - \not{p}_b}{(k_1 + k_2 - p_b)^2} \gamma_\nu \right]_{kl} \end{aligned} \quad (13)$$

where $Q_{d[u]}$ is the charge of the spectator quark.

3. Spectator corrections involving the penguin-type diagrams and the operator O_2

The corresponding diagrams are shown in Figs. 3, 4 and 5. The hard spectator contribution corresponding to the diagrams in Fig. 3a is

$$\begin{aligned} \mathcal{T}_{ijkl}^{(3a)} = & \frac{G_F}{\sqrt{2}} \frac{e}{24\pi^2} \sum_{f=u,c} V_{fd}^* V_{fb} C_2^{(f)}(\mu) \Delta F_1 \left(z_1^{(f)} \right) [\gamma_\nu]_{kl} \\ & \times \left[\left\{ \gamma_\nu - \frac{(k_2 - l)_\nu (\not{k}_2 - \not{l})}{(k_2 - l)^2} \right\} (1 - \gamma_5) \frac{\not{k}_1 + \not{k}_2 - \not{l} + m_b}{(k_1 + k_2 + l)^2 - m_b^2} \not{\epsilon}^* \right. \\ & \left. + \not{\epsilon}^* \frac{\not{p}_b + \not{l} - \not{k}_2}{(p_b + l - k_2)^2} \left\{ \gamma_\nu - \frac{(k_2 - l)_\nu (\not{k}_2 - \not{l})}{(k_2 - l)^2} \right\} (1 - \gamma_5) \right]_{ij} \end{aligned} \quad (14)$$

and from the diagrams in Fig. 3b, where the photon is emitted from the spectator quark line yield:

$$\begin{aligned}
T_{ijkl}^{(3b)} &= \frac{G_F}{\sqrt{2}} \frac{e Q_d[u]}{24\pi^2} \sum_{f=u,c} V_{fd}^* V_{fb} C_2^{(f)}(\mu) \Delta F_1 \left(z_0^{(f)} \right) \\
&\times \left[\not{\epsilon}^* \frac{k_1 + k_2 - p_b}{(k_1 + k_2 - p_b)^2} \gamma_\nu + \gamma_\nu \frac{p_b + l - k_1}{(p_b + l - k_1)^2} \not{\epsilon}^* \right]_{kl} \\
&\times \left[\left\{ \gamma_\nu - \frac{(p_b - k_1)_\nu}{(p_b - k_1)^2} (\not{p}_b - \not{k}_1) \right\} (1 - \gamma_5) \right]_{ij}
\end{aligned} \tag{15}$$

The detailed discussion about $\Delta F_1 \left(z_1^{(f)} \right)$ and $\Delta F_1 \left(z_0^{(f)} \right)$ can be found in [20].

The contributions from the diagrams in Fig. 4 can be written as

$$\begin{aligned}
T_{ijkl}^{(4)} &= -\frac{G_F}{\sqrt{2}} \frac{e}{6\pi^2} \frac{[\gamma_\nu]_{kl}}{(k_2 - l)^2 (q[k_2 - l])} \sum_{f=u,c} V_{fd}^* V_{fb} C_2^{(f)}(\mu) \\
&\times \left[\left\{ \left[\begin{aligned} &q_\nu E(k_2 - l, e^*, q) - (q[k_2 - l]) E(\nu, e^*, q) \\ &+ (e^*[k_2 - l]) E(q, \nu, k_2 - l) \\ &- (q[k_2 - l]) E(e^*, \nu, k_2 - l) \end{aligned} \right] \Delta i_5 \left(z_0^{(f)}, z_1^{(f)}, 0 \right) \right. \right. \\
&\quad \left. \left. + \left[\begin{aligned} &(k_2 - l)^2 E(\nu, e^*, q) \\ &+ (k_2 - l)_\nu E(e^*, k_2 - l, q) \end{aligned} \right] \Delta i_{25} \left(z_0^{(f)}, z_1^{(f)}, 0 \right) \right\} (1 - \gamma_5) \right]_{ij}
\end{aligned} \tag{16}$$

where

$$E(\mu, \nu, \rho) \equiv \frac{1}{2} (\gamma_\mu \gamma_\nu \gamma_\rho - \gamma_\rho \gamma_\nu \gamma_\mu) = -i \varepsilon_{\mu\nu\rho\sigma} \gamma^\sigma \gamma_5. \tag{17}$$

and the form of $\Delta i_5 \left(z_0^{(f)}, z_1^{(f)}, 0 \right)$ and $\Delta i_{25} \left(z_0^{(f)}, z_1^{(f)}, 0 \right)$ along with the detailed discussion is given in [20].

Finally, there are the diagrams where the photon is emitted from the internal quark line due to the effective $b \rightarrow s\gamma$ interaction and a gluon is exchanged between the spectator quark and the b - or s quark as shown in Fig. 5. For on shell photon such kind of diagrams do not contribute and hence the contribution comes from the Fig. 5. is zero.

Using Equations (9) and (10) along with the hard scattering matrix derived in the Eqs. (11-16), we can write from Eq. (8) as

$$\Delta \mathcal{M}_{\text{sp}}^{(K_1)} = \frac{G_F}{\sqrt{2}} \frac{e \alpha_s C_F}{4\pi N_c} f_B f_\perp^{(K_1)} M [(e^* \varepsilon^*) + i \text{eps}(e^*, \varepsilon^*, n_-, v)] \sum_{k=1}^5 \Delta H_k^{(K_1)} \tag{18}$$

where $\text{eps}(a, b, c, d) = \varepsilon_{\mu\nu\rho\sigma} a^\mu b^\nu c^\rho d^\sigma$ and the upper index K_1 characterizes the final axial meson. The dimensionless functions $\Delta H_k^{(K_1)}$ ($k = 1, 2, 3, 4, 5$) describe the contributions of the sets of Feynman diagrams presented in Figs. 1-5, respectively. In the leading order of the inverse B -meson mass ($\sim \Lambda_{\text{QCD}}/M$), the result reads as follows:

$$\Delta H_1^{(K_1)}(\mu) \simeq V_{ts}^* V_{tb} C_7^{\text{eff}}(\mu) m_b(\mu) \left[\langle l_+^{-1} \rangle_+ \langle \bar{u}^{-1} \rangle_\perp^{(K_1)}(\mu) + \langle l_+^{-1} \rangle_- \langle \bar{u}^{-2} \rangle_\perp^{(K_1)}(\mu) \right], \tag{19}$$

$$\Delta H_2^{(K_1)}(\mu) \simeq \frac{1}{3} V_{ts}^* V_{tb} C_8^{\text{eff}}(\mu) m_b(\mu) \langle l_+^{-1} \rangle_+ \langle u^{-1} \rangle_\perp^{(K_1)}(\mu), \tag{20}$$

$$\Delta H_3^{(K_1)}(\mu) \simeq 0, \tag{21}$$

$$\Delta H_4^{(K_1)}(\mu) \simeq \frac{1}{3} C_2(\mu) M \langle l_+^{-1} \rangle_+ \left[V_{ts}^* V_{tb} \langle \bar{u}^{-1} \rangle_\perp^{(K_1)}(\mu) + V_{cs}^* V_{cb} h^{(K_1)}(z, \mu) \right], \tag{22}$$

$$\Delta H_5^{(K_1)}(\mu) \simeq 0, \tag{23}$$

where $z = m_c^2/m_b^2$ and the short-hand notation used are for the integrals over the mesons distribution functions:

$$\langle l_+^N \rangle_\pm \equiv \int_0^\infty dl_+ l_+^N \phi_\pm^{(B)}(l_+), \quad \langle f \rangle_{\perp, \parallel}^{(K_1)}(\mu) \equiv \int_0^1 du f(u) \phi_{\perp, \parallel}^{(K_1)}(u, \mu), \tag{24}$$

and for convenience the following function is introduced:

$$h^{(K_1)}(z, \mu) = \left\langle \frac{\Delta i_5(z_0^{(c)}, 0, 0) + 1}{\bar{u}} \right\rangle_{\perp}^{(K_1)}. \quad (25)$$

The expressions of $\Delta H_k^{(K_1)}$ given in Eqs. (19-23) are similar to those obtained for $B \rightarrow K^* \gamma$ (c.f. Eqs. (4.4-4.6) of Ali et al. [20]) which show that the additional γ_5 present in the DA of K_1 has no effect on the calculations. Using the above Equations in Eq. (18) one can write

$$\begin{aligned} \Delta \mathcal{M}_{\text{sp}} = & \frac{G_F}{\sqrt{2}} V_{tp}^* V_{tb} \frac{\alpha_s C_F}{4\pi} \frac{e}{4\pi^2} \Delta F_{\perp}^{(K_1)}(\mu) [(pP)(e^* \varepsilon^*) + i \text{eps}(e^*, \varepsilon^*, p, P)] \\ & \times \left[C_7^{\text{eff}}(\mu) + \frac{1}{3} C_8^{\text{eff}}(\mu) \frac{\langle u^{-1} \rangle_{\perp}^{(K_1)}}{\langle \bar{u}^{-1} \rangle_{\perp}^{(K_1)}} + \frac{1}{3} C_2(\mu) \left(1 + \frac{V_{cs}^* V_{cb}}{V_{ts}^* V_{tb}} \frac{h^{(K_1)}(z, \mu)}{\langle \bar{u}^{-1} \rangle_{\perp}^{(K_1)}(\mu)} \right) \right], \end{aligned} \quad (26)$$

where

$$\Delta F_{\perp}^{(K_1)}(\mu) = \frac{8\pi^2 f_B f_{\perp}^{(K_1)}(\mu)}{N_c M \lambda_{B,+}} \langle \bar{u}^{-1} \rangle_{\perp}^{(K_1)}(\mu), \quad (27)$$

is the dimensionless quantity. $\lambda_{B,+}^{-1} = \langle l_+^{-1} \rangle_+$ is the first negative moment of the B -meson distribution function $\phi_+^{(B)}(l_+)$ which is typically estimated as $\lambda_{B,+}^{-1} = (3 \pm 1) \text{ GeV}$ [32,33]. In a recent paper by Braun et al. [34], the scale dependence of this moment is worked out at next to leading order and the value obtained is $\lambda_{B,+}^{-1}(1 \text{ GeV}) = (2.15 \pm 0.50) \text{ GeV}$. At the scale $\mu_{\text{sp}} = \sqrt{\mu_b \Lambda_H}$ of the hard-spectator corrections, and for the central values of the parameters are shown in Table I.

The analytical expression for the function $h^{(V)}(z, \mu)$ for the vector meson is given in [20]. We will proceed to give an analytical result for the axial meson. One can write the leading twist distribution amplitude $\phi_{\perp}^{(K_1)}(u, \mu)$ as [31]

$$\phi_{\perp}^{(K_1)}(u, \mu) = 6u\bar{u} \left[1 + \sum_{n=1}^{\infty} a_{\perp n}^{(K_1)}(\mu) C_n^{3/2}(u - \bar{u}) \right], \quad (28)$$

where $C_n^{3/2}(u - \bar{u})$ are the Gegenbauer polynomials [$C_1^{3/2}(u - \bar{u}) = 3(u - \bar{u})$, $C_2^{3/2}(u - \bar{u}) = 3[5(u - \bar{u})^2 - 1]/2$, etc.] and $a_{\perp n}^{(K_1)}(\mu)$ are the corresponding Gegenbauer moments. These moments are scale dependent and so should be evaluated at the scale μ ; their scale dependence is governed by [31]:

$$a_{\perp n}^{(K_1)}(\mu) = \left(\frac{\alpha_s(\mu^2)}{\alpha_s(\mu_0^2)} \right)^{\gamma_n/\beta_0} a_{\perp n}^{(K_1)}(\mu_0), \quad \gamma_n = 4C_F \left(\sum_{k=1}^n \frac{1}{k} - \frac{n}{n+1} \right), \quad (29)$$

where $\beta_0 = (11N_c - 2n_f)/3$ and γ_n is the one-loop anomalous dimension with $C_F = (N_c^2 - 1)/(2N_c) = 4/3$. In the limit $\mu \rightarrow \infty$ the Gegenbauer moments vanish, $a_{\perp n}^{(K_1)}(\mu) \rightarrow 0$, and the leading-twist transverse distribution amplitude has its asymptotic form:

$$\phi_{\perp}^{(K_1)}(u, \mu) \rightarrow \phi_{\perp}^{(\text{as})}(u) = 6u\bar{u}. \quad (30)$$

A simple model of the transverse distribution which includes contributions from the first $a_{\perp 1}^{(K_1)}(\mu)$ and the second $a_{\perp 2}^{(K_1)}(\mu)$ Gegenbauer moments only is used here in the analysis. In this approach the quantities $\langle u^{-1} \rangle_{\perp}^{(K_1)}$ and $\langle \bar{u}^{-1} \rangle_{\perp}^{(K_1)}$ are:

$$\langle u^{-1} \rangle_{\perp}^{(K_1)} = 3 \left[1 - a_{\perp 1}^{(K_1)}(\mu) + a_{\perp 2}^{(K_1)}(\mu) \right], \quad \langle \bar{u}^{-1} \rangle_{\perp}^{(K_1)} = 3 \left[1 + a_{\perp 1}^{(K_1)}(\mu) + a_{\perp 2}^{(K_1)}(\mu) \right], \quad (31)$$

and depend on the scale μ due to the coefficients $a_{\perp n}^{(K_1)}(\mu)$. The calculation for the axial K meson without Gegenbauer moments is done in detail [25,28]. In our calculation we will incorporate these effects in the calculations and will check the sensitivity of branching ratio with the LEET form factors in the presence of these moments. The Gegenbauer moments were evaluated at the scale $\mu_0 = 1 \text{ GeV}$, yielding [31]: $a_{\perp 1}^{(K^*)}(1 \text{ GeV}) = 0.20 \pm 0.05$ and $a_{\perp 2}^{(K^*)}(1 \text{ GeV}) = 0.04 \pm 0.04$ for the K^* -meson. The value of these two Gegenbauer moments have recently been modified and it has

been pointed out that these values are now larger in magnitude, have larger errors and, moreover, the first Gegenbauer moment changes its sign [35]. The new values of these Gegenbauer moments are $a_{\perp 1}^{(K^*)}(1 \text{ GeV}) = -0.34 \pm 0.18$ and $a_{\perp 2}^{(K^*)}(1 \text{ GeV}) = 0.13 \pm 0.08$ for the K^* -meson [36]. We will use the same value for the K_1 because one can see that the value is not changed for the axial meson also because changing the scale has not the noticeable effect on the coupling constants and so on the Gegenbauer moments [31], [37]. In the same manner, the function $h^{(K_1)}(z, \mu)$ introduced in Eq. (26) can be presented as an expansion on the Gegenbauer moments:

$$\begin{aligned} h^{(K_1)}(z, \mu) &= h_0(z) + a_{\perp 1}^{(K_1)}(\mu) h_1(z) + a_{\perp 2}^{(K_1)}(\mu) h_2(z) \\ &= \left[1 + 3a_{\perp 1}^{(K_1)}(\mu) + 6a_{\perp 2}^{(K_1)}(\mu) \right] \langle (\Delta i_5 + 1)/\bar{u} \rangle_{\perp}^{(0)} \\ &\quad - 6 \left[a_{\perp 1}^{(K_1)}(\mu) + 5a_{\perp 2}^{(K_1)}(\mu) \right] \langle \Delta i_5 + 1 \rangle_{\perp}^{(0)} + 30 a_{\perp 2}^{(K_1)}(\mu) \langle \bar{u} (\Delta i_5 + 1) \rangle_{\perp}^{(0)}, \end{aligned} \quad (32)$$

where another short-hand notation is introduced for the integral:

$$\langle f(u) \rangle_{\perp}^{(0)} = \int_0^1 du f(u) \phi_{\perp}^{(\text{as})}(u). \quad (33)$$

The detail of relevant functions as well as the analytical form of the $\langle (\Delta i_5 + 1)/\bar{u} \rangle_{\perp}^{(0)}$, $\langle \Delta i_5 + 1 \rangle_{\perp}^{(0)}$ and $\langle \bar{u} (\Delta i_5 + 1) \rangle_{\perp}^{(0)}$ is given [20]. The real and imaginary parts of the functions $h_n(z)$ are presented in Figs. 6 (for $n = 0$) and 7 (for $n = 1$ and $n = 2$). The dependence on $z = m_c^2/m_b^2$ of the function $h^{(K_1)}(z, \mu)$ at the mass scale $\mu = \mu_{\text{sp}} = 1.52$ GeV of hard-spectator corrections is presented in Fig. 8. We have observed that our plots given in Fig. 7 and Fig. 8 are different to those given by Ali et al. (c.f. Fig. 7 and Fig. 8) [20]. The authors of the article [20] are agree to this observation [21], pointed out by Gilani [22]. The value of the corresponding Gegenbauer moments used for the evaluation are given in Table I.

	$K_1(1270)$		$K_1(1400)$	
$\mu, [\text{GeV}]$	μ_{sp}	$m_{b,\text{pole}}$	μ_{sp}	$m_{b,\text{pole}}$
	1.52	4.65	1.52	4.65
$a_{\perp 1}(\mu)$	-0.321	-0.285	-0.321	-0.285
$a_{\perp 2}(\mu)$	0.118	0.097	0.118	0.097
$h_0(z)$	$3.91 + i1.64$	$3.91 + i1.64$	$3.91 + i1.64$	$3.91 + i1.64$
$h^{(V)}(z, \mu)$	$2.89 + i1.74$	$3.07 + i1.34$	$2.89 + i1.74$	$3.07 + i1.34$
$\langle \bar{u}^{-1} \rangle_{\perp}^{(V)}(\mu)$	2.39	2.43	2.39	2.43
$h^{(V)}/\langle \bar{u}^{-1} \rangle_{\perp}^{(V)}$	$1.21 + i0.73$	$1.26 + i0.71$	$1.21 + i0.73$	$1.26 + i0.71$
$f_{\perp}^{(V)}(\mu), [\text{MeV}]$	118.6	111.78	88.44	83.38
$\Delta F_{\perp}^{(V)}(\mu)$	0.55	0.52	0.41	0.39

The amplitude (18) is proportional to the tensor decay constant $f_{\perp}^{(K_1)}$ of the axial meson which is a scale dependent parameter. As for the Gegenbauer moments $a_{\perp n}^{(K_1)}$, there values were defined at the mass scale $\mu_0 = 1$ GeV from the LCSR is [38]: $f_{\perp}^{(K_1)}(1 \text{ GeV}) = 122 \text{ MeV}$. Their values at an arbitrary scale μ can be obtained with the help of the evolution equation [31]:

$$f_{\perp}^{(V)}(\mu) = \left(\frac{\alpha_s(\mu^2)}{\alpha_s(\mu_0^2)} \right)^{4/(3\beta_0)} f_{\perp}^{(V)}(\mu_0). \quad (34)$$

Central values of the tensor decay constants at the scales $\mu_{\text{sp}} = 1.52 \text{ GeV}$ and $m_{b,\text{pole}} = 4.65 \text{ GeV}$ are presented in Table I.

4. Branching Ratio for $B \rightarrow K_1 \gamma$

The branching ratio for $B \rightarrow K_1 \gamma$ is simply given by

$$\begin{aligned}
\mathcal{B}_{\text{th}}(B \rightarrow K^* \gamma) &= \tau_B \Gamma_{\text{th}}(B \rightarrow K^* \gamma) \\
&= \tau_B \frac{G_F^2 \alpha |V_{tb} V_{ts}^*|^2}{32\pi^4} m_{b,\text{pole}}^2 M^3 \left[\xi_{\perp}^{(K_1)} \right]^2 \left(1 - \frac{m_{K^*}^2}{M^2} \right)^3 \left| C_7^{(0)\text{eff}} + A^{(1)}(\mu) \right|^2
\end{aligned} \tag{35}$$

where G_F is the Fermi coupling constant, $\alpha = \alpha(0) = 1/137$ is the fine-structure constant, $m_{b,\text{pole}}$ is the pole b -quark mass, M and m_{K_1} are the B - and K_1 -meson masses, and τ_B is the lifetime of the B^0 - or B^+ -meson. The value of these constants is used from [20] for the numerical analysis. For this study, we consider $\xi_{\perp}^{(K_1)}$ as a free parameter and we will extract its value from the current experimental data on $B \rightarrow K_1 \gamma$ decays.

The function $A^{(1)}$ in Eq. (35) can be decomposed into the following three components:

$$A^{(1)}(\mu) = A_{C_7}^{(1)}(\mu) + A_{\text{ver}}^{(1)}(\mu) + A_{\text{sp}}^{(1)K_1}(\mu_{\text{sp}}) . \tag{36}$$

Here, $A_{C_7}^{(1)}$ and $A_{\text{ver}}^{(1)}$ are the $\mathcal{O}(\alpha_s)$ (i.e. NLO) corrections due to the Wilson coefficient C_7^{eff} and in the $b \rightarrow s \gamma$ vertex, respectively, and $A_{\text{sp}}^{(1)K_1}$ is the $\mathcal{O}(\alpha_s)$ hard-spectator corrections to the $B \rightarrow K_1 \gamma$ amplitude computed in this paper. Their explicit expressions are as follows:

$$A_{C_7}^{(1)}(\mu) = \frac{\alpha_s(\mu)}{4\pi} C_7^{(1)\text{eff}}(\mu), \tag{37}$$

$$\begin{aligned}
A_{\text{ver}}^{(1)}(\mu) &= \frac{\alpha_s(\mu)}{4\pi} \left\{ \frac{32}{81} \left[13C_2^{(0)}(\mu) + 27C_7^{(0)\text{eff}}(\mu) - 9C_8^{(0)\text{eff}}(\mu) \right] \ln \frac{m_b}{\mu} \right. \\
&\quad \left. - \frac{20}{3} C_7^{(0)\text{eff}}(\mu) + \frac{4}{27} (33 - 2\pi^2 + 6\pi i) C_8^{(0)\text{eff}}(\mu) + r_2(z) C_2^{(0)}(\mu) \right\},
\end{aligned} \tag{38}$$

$$\begin{aligned}
A_{\text{sp}}^{(1)K_1}(\mu_{\text{sp}}) &= \frac{\alpha_s(\mu_{\text{sp}})}{4\pi} \frac{2\Delta F_{\perp}^{(K_1)}(\mu_{\text{sp}})}{9\xi_{\perp}^{(K_1)}} \left\{ 3C_7^{(0)\text{eff}}(\mu_{\text{sp}}) \right. \\
&\quad \left. + C_8^{(0)\text{eff}}(\mu_{\text{sp}}) \left[1 - \frac{6a_{\perp 1}^{(K_1)}(\mu_{\text{sp}})}{\langle \bar{u}-1 \rangle_{\perp}^{(K_1)}(\mu_{\text{sp}})} \right] + C_2^{(0)}(\mu_{\text{sp}}) \left[1 - \frac{h^{(K_1)}(z, \mu_{\text{sp}})}{\langle \bar{u}-1 \rangle_{\perp}^{(K_1)}(\mu_{\text{sp}})} \right] \right\}.
\end{aligned} \tag{39}$$

Actually $C_7^{(1)\text{eff}}(\mu)$ and $A_{\text{ver}}^{(1)}(\mu)$ are process independent and encodes the QCD effects only, where as $A_{\text{sp}}^{(1)}(\mu_{\text{sp}})$ contains the key information about the out going mesons. The factor $\frac{6a_{\perp 1}^{(K_1)}(\mu_{\text{sp}})}{\langle \bar{u}-1 \rangle_{\perp}^{(K_1)}(\mu_{\text{sp}})}$ appearing in the Eq. (39) is arising due to the Gegenbauer moments. The purpose of this paper is to see the effect of these Gegenbauer moments on the value of the form factor. As it is mentioned in [20,25,28], that the non-asymptotic corrections in the K_1 meson wave-function reduces the coefficient of the anomalous chromomagnetic moment $C_8^{(0)\text{eff}}(\mu_{\text{sp}})$ by the amount 20%. Therefore it is viable to calculate the effect of these Gegenbauer moments. The value obtained for the quantity $\left| C_7^{(0)\text{eff}} + A^{(1)}(\mu) \right|^2$ at different scales is listed in the Table II for $K_1(1270)$.

m_c/m_b	0.29	0.29	0.29
μ	$\bar{m}_b = 4.27\text{GeV}$	$m_{b,\text{pole}} = 4.65\text{GeV}$	$m_{b,\text{PS}} = 4.6\text{GeV}$
$\left(C_7^{(0)\text{eff}} + A^{(1)}(\mu) \right)_{\text{Our}}$	$-0.358 - i0.022$	$-0.356 - i0.021$	$-0.356 - i0.021$
$\left(C_7^{(0)\text{eff}} + A^{(1)}(\mu) \right)_{\text{Lee}} [25]$	$-0.406 - i0.033$	$\times$	$-0.410 - i0.033$
$\left C_7^{(0)\text{eff}} + A^{(1)}(\mu) \right _{\text{Our}}^2$	0.128	0.127	0.127

Table II

The numbers given for the quantity $\left(C_7^{(0)\text{eff}} + A^{(1)}(\mu) \right)$ needs some comments. The first one is that with the same value of the quark mass ratio m_c/m_b , the total amplitude has negligible dependence on the choice of b -quark mass or in other words the scale μ . Secondly, if the effects of the Gegenbauer moments are included then one can easily see from third and fourth row of the Table II that the value of the total amplitude reduces as compared to the value given in the literature [25]. In order to calculate the numerical value for the branching ratio we use the reference scale to be

$$(\mu, \mu_{\text{sp}}) = (4.27 \text{ GeV}, 1.45 \text{ GeV})$$

After calculating the Gegenbauer moments at these scales the only independent parameter which is left in the calculation of the branching ratio is the LEET form factor and it has the biggest theoretical uncertainty.. By taking the value of the form factor from the LCSR which is $\xi_{\perp}^{(K_1)}(0) = 0.14 \pm 0.03$ it was shown some time ago that the value of the branching ratio is very small as compared to the experimental results [25]. Then the value of the form factor is extracted from the experimental measurements (2) and it is found that the value is [28]

$$\xi_{\perp}^{(K_1)}(0) = 0.32 \pm 0.03$$

which is much larger than LCSR result for the form factor. and is contrary to that of the K^* meson where the value of LEET form factor $\xi_{\perp}^{(K^*)}$ is smaller as compared to the LCSR result. Such discrepancy in case of K^* is not yet known but for the axial K meson some sources of discrepancies are discussed in [28] and it is said that it will be bad if the Gegenbauer moments increase the value of this form factor. But we have shown that it is not the case.

Now by putting the value of the total amplitude at the scale $(\mu, \mu_{\text{sp}}) = (4.27 \text{ GeV}, 1.45 \text{ GeV})$ calculated in the last row of Table II and all the other inputs from [25] in Eq. (35) one can see that the value extracted for the form factor remains the same. Thus even if we consider the non asymptotic form of the light-cone DA, it has a very small effect on the total decay amplitude and leaves all the other things almost same. The reason is that the dominant contribution comes from the operator O_7 and so from the Wilson coefficient C_7 .

In conclusion, we surveyed the implications of the first observation of $B \rightarrow K_1(1270)\gamma$ by using the higher twists in the light-cone DA that are encoded in the coefficient of Gegenbauer expansion. By incorporating all these effects the value of the relevant form factor (LEET) is extracted from the data at NLO of α_s . It is shown that the value of the form factor remains the same as calculated with out these higher twists (asymptotic form). So the non-asymptotic form of DA is not the suitable candidate to explain the discrepancy..

Acknowledgments

One of the authors (J) would like to thank Prof. Fayyazuddin for valuable discussion and also to Amjad Gilani for giving us the code [which he developed during the review of Ref. [20]] which is used to solve this problem with some modifications. This work was supported by a grant from Higher Education Commission of Pakistan.

-
- [1] See, e.g., C. Greub, talk given at the 8th International Symposium on Heavy Flavour Physics, Southampton, England, 25-29 Jul 1999. [hep-ph/9911348].
 - [2] M. S. Alam et al. [CLEO Collaboration], Phys. Rev. Lett. **74** 2885 (1995)
 - [3] S. Cheng et al. [CLEO Collaboration], Phys. Rev. Lett. **87** 251807 (2001) [hep-ex/0108032]
 - [4] R. Barate et al. [ALEPH Collaboration], Phys. Lett. **B429** 169 (1998).
 - [5] K. Chetyrkin, M. Misiak and M. Münz, Phys. Lett. **B400** 206 (1997); E: **B425** 414 (1998) [hep-ph/9612313].
 - [6] A. L. Kagan and M. Neubert, Eur. Phys. J. **C7** (1999) 5 [hep-ph/9805303].
 - [7] P. Gambino and M. Misiak, Nucl. Phys. **B611** 338 (2001) [hep-ph/0104034].
 - [8] A. Ali, E. Lunghi, C. Greub and G. Hiller, DESY 01-217 [hep-ph/0112300].
 - [9] Belle Collaboration, K. Abe et al., BELLE-CONF-0319, 2003.
 - [10] B. Aubert et al. [BaBar Collaboration], Phys. Rev. Lett. **88** 101805 (2002)
 - [11] T. E. Coan et al. [CLEO Collaboration] Phys. Rev. Lett. **84** 5283 (2000).
 - [12] J.M. Soares, Nucl. Phys. B367 (1991) 575; Phys. Rev. D **49** 283 (1994).
 - [13] C. Greub, H. Simma, and D. Wyler, Nucl. Phys. **B434** 39 (1995); Erratum *ibid.* **444** 447 (1995).
 - [14] C. Greub, T. Hurth, and D. Wyler, Phys. Rev. D **54** 3350 (1996).
 - [15] K. Adel and Y. Yao, Phys. Rev. D **49** 4945 (1994).
 - [16] K. Chetyrkin, M. Misiak, and M. Münz, Phys. Lett. B **400**, 206 (1997).
 - [17] M. Beneke and T. Feldmann, Nucl. Phys. **B592**, 3 (2001).
 - [18] M. Beneke, T. Feldmann, and D. Siedel, Nucl. Phys. **B612**, 25 (2001).
 - [19] S.W. Bosch and G. Buchalla, Nucl. Phys. **B621**, 459 (2002).
 - [20] A. Ali and A.Ya. Parkhomenko, Eur. Phys. J. C **23**, 89 (2002).
 - [21] A. Ali and A. Y. Parkhomenko, Private communication.
 - [22] A. H. S. Gilani, [hep-ph/0404026]
 - [23] J. Chay and C. Kim, Phys. Rev. D **68**, 034013 (2003).
 - [24] Belle Collaboration, K. Abe *et al.*, BELLE-CONF-0411, ICHEP04 11-0656, [arXiv:hep-ex/0408138].
 - [25] J. P. Lee, Phys. Rev. D **69** 114007 (2004) [arXiv: hep-ph/0403034].

- [26] M. Gronau, Y. Grossman, D. Pirjol, and A. Ryd, Phys. Rev. Lett. **88**, 051802 (2002); M. Gronau and D. Pirjol, Phys. Rev. D **66**, 054008 (2002).
- [27] J.-P. Lee, Phys. Rev. D **69**, 014017 (2004); *Proceeding of the 2nd ICFP 03, Seoul, Korea* [hep-ph/0312010].
- [28] Y. J. Kwon and J. P. Lee, Phys. Rev. D **71** 014009 (2005) [arXiv:hep-ph/0409133]
- [29] M. J. Dugan and B. Grinstein, Phys. Lett. B **255**, 583 (1991).
- [30] J. Charles, A. Le Yaouanc, L. Oliver, O. Pene and J. C. Raynal, Phys. Rev. D **60**, 014001 (1999) [hep-ph/9812358].
- [31] P. Ball, V. M. Braun, Y. Koike and K. Tanaka, Nucl. Phys. B **529**, 323 (1998) [hep-ph/9802299].
- [32] A. G. Grozin and M. Neubert, Phys. Rev. D **55**, 272 (1997) [hep-ph/9607366].
- [33] M. Beneke and T. Feldmann, Nucl. Phys. B **592**, 3 (2001) [hep-ph/0008255].
- [34] V. M. Braun, D. Y. Ivanov and G. P. Korchemsky, Phys. Rev. D **69**, 034014 (2004) [arXiv:hep-ph/0309330].
- [35] P. Ball and M. Boglione, Phys. Rev. D **68**, 094006 (2003) [arXiv: hep-ph/0307337].
- [36] A. Ali, E. Lunghi and A. Y. Parhomenko, Phys. Lett. B **595** (2004) 323-338 [arXiv:hep-ph/0405075].
- [37] P. Ball and V. M. Braun, Phys. Rev. D **54** 2182 (1996).
- [38] A. Salim Safir, Eur. Phys. J. direct C **3** (2001) 15 [hep-ph/0109232]

Figure Captions

a Leading order contribution by operator O_7

- 1 Feynman diagrams contributing to the spectator corrections involving the O_7 operator in the decay $B \rightarrow K_1 \gamma$.
The curly (dashed) line here and in subsequent figures represents a gluon (photon)
- 2 Feynman diagrams contributing to the spectator corrections involving the O_8 operator in the decay $B \rightarrow K_1 \gamma$.
Row a: Photon is emitted from the flavor changing quark line
Row b: Photon radiation off the spectator quark line
- 3 Feynman diagrams contributing to the spectator corrections involving the O_2 operator in the decay $B \rightarrow K_1 \gamma$.
Row a: Photon is emitted from the flavor changing quark line
Row b: Photon radiation off the spectator quark line
- 4 Feynman diagrams contributing to the spectator corrections in $B \rightarrow K_1 \gamma$ decays involving the O_2 operator for the case when both the photon and the virtual gluon are emitted from the internal (loop) quark line
- 5 Feynman diagrams contributing to the spectator corrections in $B \rightarrow K_1 \gamma$ decays involving the O_2 operator for the case when photon is emitted from the internal (loop) quark line in the $bs\gamma$ vertex.
- 6 The function $h_0(z)$ is plotted against the ratio m_f^2/m_b^2 , where m_b is the b -quark mass. The solid curve is the real part of the function and the dashed curve is the imaginary part.
- 7 The function $h^{(K_1)}(z, \mu_{\text{sp}})$ is plotted against the ratio m_f^2/m_b^2 at the mass scale of the hard spectator correction $\mu_{\text{sp}} = 1.52$ GeV. The solid curve is the real part of the function and the dashed curve is the imaginary part
- 8 The function $h_1(z)$ (left figure) and $h_2(z)$ (right figure) are plotted against the ratio m_f^2/m_b^2 , where m_b is the b -quark mass. The solid curves are the real parts of the function and the dashed curves are the imaginary parts.






Figure 1




a




b

Figure 2



a



b

Figure 3




Figure 4




Figure 5




Figure 6





Figure 7




Figure 8

